



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 5/10, A61K 48/00	A1	(11) International Publication Number: WO 00/09669 (43) International Publication Date: 24 February 2000 (24.02.00)
(21) International Application Number: PCT/US99/18403 (22) International Filing Date: 12 August 1999 (12.08.99) (30) Priority Data: 09/134,771 12 August 1998 (12.08.98) US (71) Applicant: SIGNAL PHARMACEUTICALS, INC. [US/US]; 5555 Oberlin Drive, San Diego, CA 92121 (US). (72) Inventors: SAH, Dinah, W.; 8492 Cliffridge Avenue, La Jolla, CA 92037 (US). RAYMON, Heather, K.; 535 Genter Street, La Jolla, CA 92037 (US). (74) Agents: MAKI, David, J. et al.; Seed and Berry LLP, 6300 Columbia, 701 Fifth Avenue, Seattle, WA 98104-7092 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: HUMAN MESENCEPHALON CELL LINES AND METHODS OF USE THEREFOR (57) Abstract Conditionally-immortalized human mesencephalon cell lines are provided. Such cell lines, which may be clonal, may be used to generate neurons, including dopaminergic neurons. The cell lines and/or differentiated cells may be used for the development of therapeutic agents to prevent and treat a variety of neurological diseases such as Parkinson's disease. The cell lines and/or differentiated cells may also be used in assays and for the general study of mesencephalon cell development and differentiation.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

1
Description

HUMAN MESENCEPHALON CELL LINES AND METHODS OF USE THEREFOR

5

TECHNICAL FIELD

The present invention relates generally to human mesencephalon cell lines. The invention is more particularly related to conditionally-immortalized mesencephalon cell lines capable of differentiation into dopaminergic neurons, and to differentiated cells derived from such cell lines. Such cell lines and/or differentiated cells may be used for development of therapeutic agents and for the treatment of diseases such as Parkinson's disease. The present invention is also related to the use of such cell lines and/or differentiated cells within various assays and for the study of mesencephalon development and differentiation.

15

BACKGROUND OF THE INVENTION

In the mammalian brain, dopamine systems are critical for the control of movement, hormone release, emotional balance and reward. Alteration of dopaminergic neurotransmission is involved in a variety of nervous system disorders. One such disorder is Parkinson's disease, which results from an intrastriatal deficiency in dopamine. Currently, there is no adequate method for treating or preventing the disease. L-DOPA has been administered to patients with Parkinson's disease, but such treatments are not generally considered to be effective.

Grafts of fetal neural tissue, such as mesencephalic tissue which contains dopaminergic neurons, have been shown to improve the symptoms of parkinsonism in humans and animal models. However, the use of such grafts has been limited, due in part to the difficulty in obtaining an adequate supply of transplant tissue. In addition, such tissue has not responded well to freezing, requiring the implantation of fresh tissue. In general, the use of fresh tissue is inconvenient and undesirable, since a period of time for evaluation of the tissue (*e.g.*, to identify any viral contaminants) is beneficial.

Accordingly, there is a need in the art for stable mesencephalon cell lines that can be readily differentiated, and can be used as a renewable source of human mesencephalon dopaminergic neurons for Parkinson's research and drug discovery. The present invention fulfills these needs and further provides other related advantages.

SUMMARY OF THE INVENTION

Briefly stated, the present invention provides conditionally-immortalized human mesencephalon cell lines capable of differentiation into neurons. In one aspect, the present invention provides methods for producing a conditionally-immortalized human mesencephalon neural precursor cell, comprising the steps of: (a) transfecting human mesencephalon cells plated on a first surface and in a first growth medium that permit proliferation with DNA encoding a selectable marker and an externally regulatable growth-promoting gene; and (b) selecting the transfected cells on a second surface and in a second growth medium that permit attachment and proliferation, and therefrom producing a conditionally-immortalized human mesencephalon neural precursor cell. Within certain embodiments, the first and second surfaces are independently selected from the group consisting of substrates comprising one or more of a polyamino acid (*e.g.*, polylysine or polyornithine), fibronectin, laminin or tissue culture plastic. The growth-promoting gene may be an oncogene, such as v-myc, and expression of the growth-promoting gene may, but need not, be inhibited by tetracycline.

Within other aspects, the present invention provides conditionally-immortalized human mesencephalon neural precursor cells capable of differentiation into dopaminergic neurons and conditionally-immortalized human mesencephalon neural precursor cells capable of differentiation into GABA-ergic neurons.

The present invention further provides methods for producing neurons, comprising culturing a conditionally-immortalized human mesencephalon neural precursor cell as described above under conditions inhibiting expression of the growth-promoting gene. Within certain embodiments, the cell may be cultured in medium comprising tetracycline and/or in the presence of one or more differentiating agents such as forskolin, GDNF, CNTF, IGF-I and/or BDNF.

Within further aspects, the present invention provides neurons produced as described above.

The present invention further provides, within other aspects, methods for transplanting a human mesencephalon cell into a mammal, comprising administering to a mammal a cell produced as described above.

Within further aspects, methods are provided for treating Parkinson's disease in a patient, comprising administering to a patient a cell produced as described above.

In other aspects, methods are provided for screening for an agent that modulates an activity of a protein produced by a human mesencephalon cell, comprising: (a) contacting a cell produced as described above with a candidate agent; and (b) subsequently measuring the ability of the candidate agent to modulate an activity of a protein produced by the cell.

Within further aspects, the present invention provides methods for detecting the presence or absence of a protein in a sample, comprising: (a) contacting a sample with a cell produced as described above; and (b) subsequently detecting a response in the cell, and therefrom detecting the presence of a protein in the sample.

The present invention further provides methods for identifying a human mesencephalon gene or protein, comprising detecting the presence of a gene or protein within a culture of cells produced as described above.

In further aspects, methods are provided for screening for an agent that affects human mesencephalon cell death, comprising: (a) contacting a cell produced as described above with a candidate agent under conditions that, in the absence of candidate agent, result in death of the cell; and (b) subsequently measuring the ability of the candidate agent to affect the death of the cell.

The present invention further provides methods for screening for a protein that regulates human mesencephalon cell death, comprising: (a) altering the level of expression of a protein within a cell produced as described above; and (b) subsequently measuring the effect of the alteration on the death of the cell, and therefrom identifying a protein that regulates human mesencephalon neural precursor cell death.

Within other aspects, the present invention provides conditionally-immortalized human mesencephalon neural precursor cells produced as described above. Such cells may be present within a clonal cell line.

These and other aspects of the present invention will become apparent upon reference to the following detailed description and attached drawings. All references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A and 1B are phase contrast micrographs of immortalized human mesencephalon cells. Figure 1A shows the cells after growth in proliferative conditions. Figure 1B shows the cells after 6 days of differentiation with DMEM/F12 medium containing
5 N2 supplements, forskolin (10 μ M), BDNF (20 ng/mL) and GDNF (20 ng/mL).

Figures 2A-2D are phase contrast micrographs (Figures 2A and 2C) and photographs (Figures 2B and 2D) showing labeling for v-myc in representative immortalized human mesencephalon cells after growth in the proliferative growth condition (Figures 2A and 2B) or after differentiation (Figures 2C and 2D) for 6 days. For each field, a phase
10 contrast photograph (post-fixation) is shown to the left of the corresponding immunofluorescence photograph.

Figure 3 is a photograph showing immunofluorescence labeling for tyrosine hydroxylase (TH) in a representative immortalized human mesencephalon cell line after 6 days of differentiation. The culture was differentiated with DMEM/F12 medium containing
15 N2 supplements, tetracycline (1 μ g/mL), forskolin (10 μ M), BDNF (20 ng/mL), GDNF (20 ng/mL), CNTF (20 ng/mL) and IGF-I (100 ng/mL).

Figure 4 is a photograph showing immunofluorescence labeling for GABA in a representative immortalized human mesencephalon cell line after 6 days of differentiation. The culture was differentiated with DMEM/F12 medium containing N2 supplements,
20 tetracycline (1 μ g/mL), forskolin (10 μ M), BDNF (20 ng/mL), GDNF (20 ng/mL), CNTF (20 ng/mL) and IGF-I (100 ng/mL).

DETAILED DESCRIPTION OF THE INVENTION

As noted above, the present invention is generally directed to conditionally-
25 immortalized human mesencephalon cell lines, differentiated cells generated from such cell lines and various methods employing such cells. In particular, the present invention is directed to conditionally-immortalized human mesencephalon neural precursor cells that are capable of differentiation into dopaminergic neurons, and to the use of such cells for drug discovery and development, transplantation studies, therapeutic methods and a variety of
30 assays. Conditionally-immortalized human mesencephalon neural precursor cell lines of the present invention may, but need not, be clonal cell lines. The cell lines described herein

provide an infinite, renewable supply of homogeneous cells and facilitate treatment of, and drug development for, disorders such as Parkinson's disease.

Conditionally-immortalized human mesencephalon neural precursor cells may generally be prepared from human mesencephalon tissue (*e.g.*, fetal mesencephalon tissue).

5 Such tissue is preferably dissociated, using standard procedures. The tissue is then washed and plated on a surface and in a growth medium that permits proliferation (*i.e.*, the surface and medium permit at least about 1% of the cells to double in a 24 hour period). One preferred growth medium contains DMEM/F-12, 10% fetal calf serum and FGF-2 (human recombinant, 40 ng/mL, Boehringer Mannheim, Indianapolis, IN). Suitable surfaces include,
10 but are not limited to, one or a combination of polyamino acids (*e.g.*, polylysine and/or polyornithine), tissue culture plastic and surfaces treated with laminin or fibronectin. Cells may generally be plated at a density ranging from 10^3 to 10^5 cells/cm², preferably at a density of approximately 3×10^4 cells/cm².

Human mesencephalon neural precursor cells may be conditionally
15 immortalized by transfection of the plated cells with any suitable vector containing a growth-promoting gene (*i.e.*, a gene encoding a protein that, under appropriate conditions, promotes growth of the transfected cell) such that the production and/or activity of the growth-promoting protein is regulatable by an external factor. In a preferred embodiment the growth-promoting gene is an oncogene such as, but not limited to, *v-myc*. Other oncogenes that may
20 be used as growth promoting genes include *N-myc*, *c-myc*, p53, SV40 large T antigen, polyoma large T antigen, E1a adenovirus and E7 protein of human papillomavirus. In general, a "growth promoting gene" is any gene that, when employed within a tet-controlled expression system as described herein, results in the generation of cultures of neural precursor cells which can be differentiated into neurons.

25 External regulation of the growth-promoting protein may be achieved by placing the growth-promoting gene under the control of an externally-regulatable promoter (*i.e.*, a promoter whose activity may be controlled by, for example, modifying the temperature of the transfected cells or the composition of the medium in contact with the cells). In general, regulation of expression of the growth-promoting gene should be relatively tight (*i.e.*,
30 expression of the growth-promoting gene should generally be undetectable by immunofluorescent techniques as described herein when the promoter is repressed. For example, a tetracycline (tet)-controlled gene expression system may be employed (*see* Gossen

et al., *Proc. Natl. Acad. Sci. USA* 89:5547-5551, 1992; Hoshimaru et al., *Proc. Natl. Acad. Sci. USA* 93:1518-1523, 1996). In the absence of tet, a tet-controlled transactivator (tTA) within this vector strongly activates transcription from ph_{CMV}^{*-1} , a minimal promoter from human cytomegalovirus fused to tet operator sequences. tTA is a fusion protein of the repressor (tetR) of the transposon-10-derived tet resistance operon of *E. coli* and the acidic domain of VP16 of herpes simplex virus. Low, non-toxic concentrations of tet (0.01-1.0 $\mu\text{g/mL}$) almost completely abolish transactivation by tTA (*i.e.*, *v-myc* is no longer detectable using an immunofluorescence assay as provided herein).

In a preferred embodiment, the vector further contains a gene encoding a selectable marker (*e.g.*, a protein that confers drug resistance). The bacterial neomycin resistance gene (neo^R) is one such marker that may be employed within the present invention. Cells carrying neo^R may be selected by means known to those of ordinary skill in the art, such as the addition of 25-200 $\mu\text{g/mL}$ G418 to the growth medium. It will be readily apparent that other markers may be employed, and appropriate selections may be readily performed by those of ordinary skill in the art.

Transfection may be achieved by any of a variety of means known to those of ordinary skill in the art including, but not limited to, retroviral infection. In general, plated cells may be transfected by infection with a suitable retrovirus (*e.g.*, VSV-G pseudotyped LINX *v-myc* retrovirus, as described further below). The use of VSV-G pseudotyped retrovirus is preferred, in order to obtain higher stock concentrations of virus, to obtain stocks in the medium of choice (after centrifugation) and to increase the infectivity of human cells. Recently developed (nontraditional) VSV-G pseudotyped retroviral vectors may be especially useful for the infection of human cells, since the receptor for the VSV-glycoprotein is more abundant and less species-specific than the receptors for traditional amphotropic envelope proteins. Moreover, VSV-G pseudotyped viral particles have been reported to withstand ultracentrifugation, allowing concentration of virus and resuspension in growth medium compatible with neural progenitor cell growth (Burns et al., *Proc. Natl. Acad. Sci. USA* 90:8033-8037, 1993).

For example, a mesencephalon neural precursor cell culture prepared as described above may be infected within five days after plating by incubation for about 12-24 hours (*e.g.*, overnight) with retrovirus in the presence of polybrene (4-8 $\mu\text{g/mL}$). Retrovirus

may then be removed by washing with fresh growth medium. Transfected cells carrying a selectable marker may generally be selected on a surface and in a growth medium that permit attachment and proliferation. The ability of a surface to permit attachment may be determined using visual microscopic inspection. In general, at least about 20% of the cells should adhere to the surface. One preferred growth medium contains DMEM/F-12 with N2 supplement (GIBCO, Baltimore, MD), conditioned medium from rat CNS progenitor cells (50%; prepared as described by Sah et al., *Nature Biotechnology* 15:574-580, 1997), FGF-2 (human recombinant, 40 ng/mL, Boehringer Mannheim, Indianapolis, IN), EGF (human recombinant, 40 ng/mL, GIBCO, Baltimore, MD) and PDGFA/B (human recombinant platelet-derived growth factor, 20 ng/mL, Boehringer Mannheim, Indianapolis, IN). Suitable surfaces include, but are not limited to, one or a combination of a polyamino acid (e.g., polylysine and/or polyornithine), tissue culture plastic and surfaces treated with laminin or fibronectin, as described above, and should grow as an adherent monolayer.

Following transfection, cultures may be maintained in a simplified growth medium containing, for example, DMEM/F-12 with N2 supplement, FGF-2 (40 ng/mL), EGF (40 ng/mL) and PDGFA/B (20 ng/mL). Cultures approaching confluence may be passaged by trypsinization and split 1:5. Typically, one T75 flask near confluence may yield 10^7 cells, and cultures may be passaged every 3 to 7 days. Cells may also be frozen in liquid nitrogen for long-term storage.

Clonal cell lines may be isolated from a conditionally-immortalized human mesencephalon neural precursor cell line prepared as described above. In general, such clonal cell lines may be isolated using standard techniques, such as by limit dilution or using cloning rings, and expanded. Clonal cell lines may generally be fed and passaged as described above. Genomic Southern blots may be performed to confirm clonality.

Conditionally-immortalized human mesencephalon neural precursor cell lines (which may, but need not, be clonal) may generally be induced to differentiate into neurons by inhibiting the expression of the growth-promoting gene (i.e., suppressing the production and/or activity of the growth-promoting protein). For example, if the gene encoding the growth-promoting protein is under the control of an externally-regulatable promoter, the conditions (e.g., temperature or composition of medium) may be modified to suppress transcription of the growth-promoting gene. For the tetracycline-controlled gene expression system discussed above, differentiation may be achieved by the addition of tetracycline to

suppress transcription of the growth-promoting gene. In general, 1-5 $\mu\text{g/mL}$ tetracycline for 48 hours is sufficient to begin neuronal morphological differentiation, and the number of differentiated neurons increases during subsequent days. Such differentiation may be performed, for example, by plating the cells on a suitable substrate (*e.g.*, one or a combination of polyamino acid, fibronectin, laminin or tissue culture plastic) in a medium consisting of DMEM/F-12 with N2 supplement and tetracycline (1-5 $\mu\text{g/mL}$). It has been found, within the context of the present invention, that differentiation is enhanced by the addition of forskolin (10 μM), GDNF (glial cell-derived neurotrophic factor; 20 ng/mL), CNTF (ciliary neurotrophic factor; 20 ng/mL), IGF-I (insulin-like growth factor; 100 ng/mL) and BDNF (brain-derived neurotrophic factor; 20 ng/mL). Culture medium may then be refreshed (*e.g.*, every 2-4 days).

Certain differentiated cells are dopaminergic neural cells (*i.e.*, neural cells that express dopamine). Such cells may be identified based on the presence of tyrosine hydroxylase (TH), an enzyme that is involved in the synthesis of dopamine. In general, a cell that expresses TH at a level detectable using standard immunofluorescence techniques is considered dopaminergic. GABA-ergic differentiated cells may be similarly identified based on the detection of GABA via immunofluorescence.

Characterization of both progenitor and differentiated cell lines may generally be performed using techniques well known to those of ordinary skill in the art, including morphological analysis of cell type, immunocytochemistry and PCR (to identify cell type-specific markers and receptors and to confirm the presence of the growth-promoting gene) and electrophysiological analysis of voltage- and ligand-gated currents. Briefly, neuronal cells may be identified morphologically based on the presence of phase bright cell bodies and long, thin processes. As noted above, neuronal markers include TH, which is a marker for dopaminergic neurons, and GABA, which is a marker for GABA-ergic neurons. Map2ab may also be used as a pan-neuronal marker. The presence or absence of such markers may be readily determined using standard immunofluorescence techniques (employing, for example, commercially available primary antibodies and fluorescent reagents) and the levels of mRNA encoding such markers may be determined using PCR or hybridization techniques. Electrophysiological analyses familiar to those of ordinary skill in the art may be employed to evaluate the ability of the cells to fire action potentials and to express sodium, calcium and potassium currents, as well as ligand-gated currents (*e.g.*, dopamine (DA), N-methyl-D-

aspartate (NMDA), kainate (KA) and γ -amino-n⁹-butyric acid (GABA)), thereby determining the levels of functional channels and receptors.

Human conditionally-immortalized mesencephalon neural precursor cells may generally be used to produce dopamine-producing neurons *in vitro* or *in vivo*. For *in vivo* use, 5 conditionally-immortalized mesencephalon neural precursor cells may be implanted, such that the cells differentiate *in vivo*. Alternatively, differentiated dopaminergic neurons generated *in vitro* may be implanted.

Implantation may generally be performed using any suitable technique known in the art. For example, cells may be inserted into a delivery device (such as a syringe) that 10 facilitates introduction (by, for example, injection) of the cells to a desired location. Cells are generally present within a pharmaceutical composition as described herein for implantation.

Cells can generally be assessed for their ability to treat Parkinson's disease using any of a variety of animal models of the disease. Suitable rat models include rats having lesions in substantia nigral dopaminergic cells caused by treatment with 6- 15 hydroxydopamine, 1 methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) or by surgical transection of the nigral striatal pathway (*see, e.g.,* Bjorklund et al., *Nature* 298:652-654, 1982). A rhesus monkey or sheep model may also be used, in which the animals have lesions in substantia nigral dopaminergic cells caused by treatment with MPTP (*see, e.g.,* Smith et al., *Neuroscience* 52:7-16, 1993; Bakay et al., *Appl. Neurophysiol.* 48:358-361; Zamir et al., 20 *Brain Res.* 322:356-360, 1984; and Baskin et al., *Life Sci.* 54:471-479, 1994). Cells may be implanted using standard techniques and morphological and immunohistochemical studies can be performed to determine whether the implant has integrated into the surrounding tissue. Behavioral tests, such as the rotational symmetry model (Freed et al., *in* Sladek et al. (eds.) *Neural Transplants: Development and Function* (Plenum Press, NY):373-406, 1984), may be 25 used to confirm functional integration.

For treatment of a patient, conditionally-immortalized human mesencephalon neural precursor cell lines and/or modulating agents (*e.g.,* which may inhibit or enhance an activity of a protein produced by a human mesencephalon neural precursor cell, or may inhibit the death of differentiated human mesencephalon neural precursor cells, as described 30 below) may be administered to a patient (either prophylactically or for treatment of an existing condition). Conditions that may be prevented and/or treated using such cells and/or modulating agents include, but are not limited to, Parkinson's disease. In particular, for

treatment of Parkinson's disease, differentiated dopaminergic neural cells may be transplanted into, for example, the striatum of a patient. Cells may generally be introduced into the patient by grafting, using standard techniques. Modulating agents may be administered by any of a variety of routes known to those of ordinary skill in the art. Such agents may be administered in their active form, as prodrugs (*i.e.*, compounds that are converted to the active form within the patient) or as nucleic acid sequences encoding the modulating agent or prodrug. Conditionally-immortalized human mesencephalon neural precursor cells for use in this aspect of the present invention may, but need not, be further transfected such that they express one or more additional proteins (such as modulating agents) within the patient.

For administration to a patient, one or more conditionally-immortalized human mesencephalon neural precursor cells (and/or modulating agents) are generally formulated as a pharmaceutical composition. A pharmaceutical composition may be a sterile aqueous or non-aqueous solution, suspension or emulsion, which additionally comprises a physiologically acceptable carrier (*i.e.*, a non-toxic material that does not interfere with the activity of the active ingredient). Any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of the present invention. The selection of a carrier will depend, in part, on the nature of the substance (*i.e.*, cells or chemical compounds) being administered. Representative carriers include physiological saline solutions, gelatin, water, alcohols, natural or synthetic oils, saccharide solutions, glycols, injectable organic esters such as ethyl oleate or a combination of such materials. Optionally, a pharmaceutical composition may additionally contain preservatives and/or other additives such as, for example, antimicrobial agents, anti-oxidants, chelating agents and/or inert gases, and/or other active ingredients.

Routes and frequency of administration, as well as doses, will vary from patient to patient and with the nature of the substance being administered. In general, the pharmaceutical compositions comprising modulating agents may be administered intravenously, intramuscularly, subcutaneously or intracavity. Doses may be administered daily. Compositions comprising cells are generally implanted one or more times. A suitable dose is an amount that is sufficient to show improvement in the symptoms of a patient afflicted with a neurological condition such as Parkinson's disease or to inhibit the onset of such a condition. Symptom improvement may be detected based on improvement and/or delay in clinical symptoms associated with the disease. Symptom improvement may, but

need not, occur weeks or months after administration or implantation. In general, the amount of modulating agent present in a dose, or produced *in situ* by DNA present in a dose, ranges from about 1 mg to about 200 mg per kg of host. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.1 mL to about 5mL for 10-60 kg animal.

5 Within certain aspects of the present invention, conditionally-immortalized human mesencephalon neural precursor cell lines may be used in a variety of *in vitro* assays and screens. It will be apparent that the cell lines described herein may be used within numerous well known assays and screens, and that specific parameters and criteria for performing these methods will depend upon the assay being performed. Those of ordinary
10 skill in the art can readily design specific assays and screens based upon well known methods and upon the desired properties of the compounds to be identified.

 Within certain assays, the differentiated or undifferentiated conditionally-immortalized human mesencephalon neural precursor cell lines described herein may be used to detect the presence or absence of a nucleic acid molecule or protein of interest in such
15 cells. To detect a particular nucleic acid sequence (*i.e.*, DNA and/or RNA) within such cells, the well known methods of PCR and/or various hybridization techniques may be employed. Such assays may be readily designed and performed using methods described, for example, in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989. To detect a protein, the detection reagent is
20 typically an antibody, which may be prepared as described below. There are a variety of assay formats known to those of ordinary skill in the art for using an antibody to detect a protein in a sample. *See, e.g.*, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. Antibodies for use in such assays may be polyclonal or monoclonal. Antibodies may be prepared by any of a variety of techniques known to those of
25 ordinary skill in the art and monoclonal antibodies specific for a protein of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.* 6:511-519, 1976, and improvements thereto. Alternatively, a protein may be detected based upon its activity, using any suitable assay known in the art.

 Such assays may generally be used, for example, within methods to evaluate
30 the ability of an agent to modulate an activity of a protein produced by a human mesencephalon neural precursor cell or a neuron. Within such assays, differentiated or undifferentiated cells may be contacted with a candidate agent under conditions and for a

time sufficient to permit an agent to modulate protein activity. Following contact, the ability of the candidate agent to modulate an activity of a protein produced by the cell is measured using standard techniques, such as PCR or hybridization (for evaluating levels of mRNA) or any of a variety of immunoassays or functional assays appropriate for the protein of interest.

5 For example, calcium-sensitive or voltage-sensitive dye coupled assays, cAMP measurements and/or receptor binding assays may be employed to evaluate the effect of a candidate modulating agent. In general, a suitable amount of antibody or other agent for use in such a screen will vary depending on the particular protein, but will range from about 10 μ g to about 100 mg.

10 The term "modulation," as used herein, includes the suppression or enhancement of an activity of a protein of interest. Such modulation may occur at the transcriptional or translational level, or may be the result of altering the activity of the intact protein. Modulation of protein activity may be direct (*i.e.*, the modulating agent may interact directly with the protein of interest) or may be indirect (*i.e.*, the modulating agent may alter
15 the expression and/or activity of one or more other proteins which in turn modulate the activity of the protein of interest). Modulating agents may be antibodies (*e.g.*, monoclonal), polynucleotides or other drugs. Agents that modulate the activity of any cellular protein may be identified within such screens. In certain embodiments, modulating agents may be identified for proteins such as neurotransmitter receptors (*e.g.*, DA-receptors, AMPA-
20 preferring receptors, kainate receptors, GABA receptors, adenosine receptors and/or 5-HT receptors), growth factor receptors (*e.g.*, receptors for FGF-2, EGF, BDNF, NGF, CNTF, NT-3 and/or GDNF) or ion channels (*e.g.*, sodium channels, calcium channels and/or potassium channels). Preferred modulating agents are able to suppress or enhance the activity of a protein by at least a factor of 2.

25 In another aspect of the present invention, the cell lines described herein may be used within a system for studying protein and/or gene expression in a native neural precursor or neuronal cellular environment. For example, receptor expression and/or activity may be assayed, and the effect of various modifications on such expression and/or activity may be evaluated using methods well known to those of ordinary skill in the art. In one such
30 method, cell lines may be permanently or transiently transfected with one or more genes of interest such as, but not limited to, genes that produce or modify membrane proteins, secreted proteins or intracellular proteins of interest. Such genes include ion channels,

neurotransmitter receptors, proteins mutated in familial forms of neurodegenerative diseases (e.g., synuclein) and/or MAP kinases. The transfected genes may also be coupled to reporter genes for use in drug development. Within this and other aspects described herein, conditionally-immortalized human mesencephalon neural precursor cells may be employed
5 without differentiation, or differentiated cells may be used. In addition, cells of varying ages and grown in any of a variety of conditions may be employed. The cell lines of the present invention have many advantages over existing cell lines for such studies, including the ability to provide clonal cell lines capable of producing neurons, and the property of conditional-immortalization, which allows arrest at specific stages of development. The selection of
10 particular cells for any given study will depend on the goals of the study, and those of ordinary skill in the art will be readily able to prepare appropriate cells using the techniques described herein.

Within further aspects, conditionally-immortalized human mesencephalon neural precursor cell lines of the present invention may be used in assays to detect the
15 presence or absence of a particular protein in a sample. In general, an assay may be performed by contacting such cells with a sample and then measuring a response induced by the protein within the cells using methods familiar to those of ordinary skill in the art. For example, a response may be measured using differential display techniques.

In a further aspect, the conditionally-immortalized human mesencephalon
20 neural precursor cell lines described herein may be used in the identification of novel genes and proteins present in proliferative and differentiated (e.g., dopaminergic neuronal) human mesencephalon neural precursor cells. Techniques, such as PCR, differential display, hybridization, expression library screens, immunoassays and two-hybrid screens may be employed for such identification. A particularly useful technique is differential gene
25 screening. Clonal cell lines as described herein are particularly suited to such studies because they are derived from a single parental cell and, therefore, human mesencephalon neural precursor cell-specific genes are amplified with respect to non-clonal cell lines. Novel genes and proteins that are expressed upon experimental manipulation (e.g., induction of apoptosis) may also be identified.

30 The cell lines provided herein may also be used in *in vitro* models of neuronal cell death including, but not limited to, neuronal apoptosis induced by growth factor withdrawal. Briefly, a clonal conditionally-immortalized human mesencephalon neural

precursor cell line may be differentiated under conditions designed to minimize the basal level of apoptosis. Suitable conditions may be readily identified by evaluating the percentage of apoptotic nuclei in cells grown under different test conditions. The percentage of apoptotic nuclei may generally be determined by methods well known to those of ordinary skill in the art, such as by DAPI staining or the *in situ* nick end-labeling assay. Suitable conditions for minimizing basal apoptosis include differentiation in the presence of 1 μ g/mL tetracycline, forskolin (10 μ M), IGF-I (100 ng/mL), CNTF (20 ng/mL), GDNF (20 ng/mL) and BDNF (20 ng/mL). Cells should be maintained in suitable differentiation conditions for a time sufficient to allow differentiation, while minimizing the basal level of apoptosis (which generally increases during the first 10 days of differentiation).

Such differentiated neuronal cells may then be employed in any of several models of apoptosis. In one such model, growth factors and N2 supplement are withdrawn for an amount of time sufficient to significantly increase the percentage of apoptotic cells with condensed nuclei. Preferably, the percentage of such apoptotic cells increases by at least about two fold. Under the representative conditions described above, about 18 hours of withdrawal is generally sufficient. After 48 hours of withdrawal, a significant percentage of the cells should be rendered nonviable. Preferably, at least about 50% of the neurons are no longer viable following withdrawal of growth factors.

Regardless of the particular model, the cells may be used to study the mechanisms of apoptosis, as well as the effect of various conditions and agents on the apoptosis of neuronal cells, using experimental techniques well known to those of ordinary skill in the art. For example, the cells may be used to screen for an agent that affects neuronal cell death. Such a screen may be performed by contacting the cells during growth factor withdrawal with a candidate agent and then evaluating the ability of the candidate agent to affect the subsequent level of apoptosis. Similarly, the cells may be used to screen for a protein that regulates neuronal cell death. In such a screen, the level of expression or activity of a candidate protein (*e.g.*, an enzyme) is altered within the cells (using standard techniques) and then the affect of the alteration on the level of apoptosis following treatments (including, but not limited to, growth factor withdrawal) is measured.

The following Example is offered by way of illustration and not by way of limitation.

15
EXAMPLE

Preparation of Human Mesencephalon Progenitor Cell Lines

This Example illustrates the conditional immortalization of human mesencephalon progenitor cells.

5 Human mesencephalon cells in primary culture were prepared from first trimester human fetal brain (obtained through Advanced Bioscience Resources, Inc.). The tissue was procured in compliance with state and federal laws and regulations. The mesencephalon was dissociated by incubation at 37°C for 30 to 45 minutes in an enzyme solution containing protease 23 (3 mg/mL), with occasional trituration. The dispersed
10 mesencephalon cells were washed with saline containing trypsin inhibitor (1 mg/mL) and bovine serum albumin (1 mg/mL). The cells were subsequently plated at a density of approximately 3×10^4 cells/cm² in growth medium consisting of DMEM/F-12, 10% fetal calf serum and FGF-2 (human recombinant, 40 ng/mL, Boehringer Mannheim, Indianapolis, IN).

For retroviral infection, the LINX *v-myc* vector was employed (Hoshimaru et
15 al., *Proc. Natl. Acad. Sci. USA* 93:1518-1523, 1996; Sah et al., *Nature Biotechnol.* 15:574-580, 1997). In this system, in the absence of tetracycline, a tetracycline-controlled transactivator (tTA) strongly activates transcription from $ph_{CMV^{+}}$, resulting in the expression of the downstream *v-myc* oncogene. Tetracycline (0.01-1.0 µg/mL) almost completely abolishes transcriptional activation by tTA, thereby blocking transcription of the *v-myc*
20 oncogene. A gene conferring neomycin resistance is also present in the vector. Mesencephalon cultures were retrovirally infected and G418- selected using methods analogous to those described by Sah et al., *Nature Biotechnol.* 15:574-580, 1997.

After G418 selection, cultures were maintained in a simplified growth medium consisting of DMEM/F-12 with N2 supplement, FGF-2 (40 ng/mL), EGF (40 ng/mL) and
25 PDGF A/B (20 ng/mL). Cultures approaching confluence were passaged by trypsinization and split 1:5. Typically, one T75 flask near confluence yielded 10^7 cells, and cultures were passaged every 3 to 7 days.

During G418 selection, some cell death occurred; after the selection, *v-myc*⁺ cells became the predominant cell type in the culture. These G418- resistant, *v-myc*⁺ cells
30 grew as an adherent monolayer. The majority of the cells were polygonal with very short processes.

Differentiation of cell lines with tetracycline resulted in increased neuronal differentiation (Figure 1) as well as suppression of v-myc oncoprotein expression (Figures 2A-2D). An increase in TH expression was also observed for some cell lines, indicating the presence of dopaminergic neurons.

5 Clonal cell lines were isolated by limit dilution in 96-well plates. Single colonies were expanded by feeding and passaging them as described above. Of eighteen cultures expanded, fourteen appear to be derived from a single cell.

 One clone (MESII(1)-C2) was differentiated in DMEM/F-12 with N2 supplement, tetracycline (1 μ g/mL), forskolin (10 μ M), GDNF (20 ng/mL) and BDNF (20
10 ng/mL). Following differentiation, a high number of MAP2ab and TH immunoreactive cells were observed. For such immunocytochemistry, cells were fixed with 4% paraformaldehyde, and were incubated with primary antibody in blocking buffer for 2 hours at room temperature, rinsed, and then incubated with a fluorescein (FITC)- or Texas red-conjugated species-specific secondary antibody (Jackson ImmunoResearch Laboratories, Inc., West Grove, PA) in
15 blocking buffer for another hour at room temperature. Cultures were then rinsed three times with PBS and coverslipped with PVA/DABCO before scoring and photographing representative fields.

 Clone MESII(1)-C2 was also differentiated as described above, but with the addition of CNTF (20 ng/mL) and IGF-1 (100 ng/mL). Approximately 11% of the
20 differentiated cells were MAP2ab immunoreactive and 5% of the cells were TH immunoreactive (Figure 3). In these differentiation conditions, 41% of the MAP2ab positive cells were also TH positive and 100% of the TH positive cells were also MAP2ab positive.

 These results indicate that these immortalized human mesencephalon cell lines differentiate into dopaminergic neurons. In addition, other immortalized human
25 mesencephalon cell lines differentiate into GABA-ergic neurons (Figure 4).

 From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for the purpose of illustration, various modifications may be made without deviating from the spirit and scope of the invention.
30 Accordingly, the invention is not limited except by the appended claims.

Claims

1. A method for producing a conditionally-immortalized human mesencephalon neural precursor cell, comprising:
 - (a) transfecting human mesencephalon cells plated on a first surface and in a first growth medium that permit proliferation with DNA encoding a selectable marker and an externally regulatable growth-promoting gene; and
 - (b) selecting the transfected cells on a second surface and in a second growth medium that permit attachment and proliferation, and therefrom producing a conditionally-immortalized human mesencephalon cell.
2. The method of claim 1 wherein the first and second surfaces are independently selected from the group consisting of substrates comprising one or more of a polyamino acid, fibronectin, laminin or tissue culture plastic.
3. The method of claim 1 wherein the growth-promoting gene is an oncogene.
4. The method of claim 3 wherein the oncogene is v-myc.
5. The method of claim 1 wherein expression of the growth-promoting gene is inhibited by tetracycline.
6. A conditionally-immortalized human mesencephalon neural precursor cell capable of differentiation into neurons.
7. A conditionally-immortalized human mesencephalon neural precursor cell according to claim 6, wherein the cell is capable of differentiation into dopaminergic neurons.
8. A conditionally-immortalized human mesencephalon neural precursor cell according to claim 6, wherein the cell is capable of differentiation into GABA-ergic neurons.
9. A method for producing neurons, comprising culturing a cell produced according to claim 1 under conditions inhibiting expression of the growth-promoting gene.

10. A method according to claim 9, wherein the cell is cultured in medium comprising tetracycline.
11. A method according to claim 9, wherein the cells are cultured in the presence of one or more differentiating agents.
12. A method according to claim 11, wherein the differentiating agent is selected from the group consisting of forskolin, GDNF, CTNF, IGF-I and BDNF.
13. A neuron produced according to the method of claim 9.
14. A dopaminergic neuron produced according to the method of claim 9.
15. A GABA-ergic neuron produced according to the method of claim 9.
16. A method for transplanting a human mesencephalon cell into a mammal, comprising administering to a mammal a cell produced according to the method of claim 1 or claim 9.
17. A method for treating Parkinson's disease in a patient, comprising administering to a patient a cell produced according to the method of claim 1 or claim 9.
18. A method for screening for an agent that modulates an activity of a protein produced by a human mesencephalon cell, comprising:
 - (a) contacting a cell produced according to the method of claim 1 or claim 9 with a candidate agent; and
 - (b) subsequently measuring the ability of the candidate agent to modulate an activity of a protein produced by the cell.
19. A method for detecting the presence or absence of a protein in a sample, comprising:
 - (a) contacting a sample with a cell produced according to the method of claim 1 or claim 9; and
 - (b) subsequently detecting a response in the cell, and therefrom detecting the presence of a protein in the sample.

20. A method for identifying a human mesencephalon gene or protein, comprising detecting the presence of a gene or protein within a culture of cells produced according to the method of claim 1 or claim 9.

21. A method for screening for an agent that affects human mesencephalon cell death, comprising:

(a) contacting a cell produced according to the method of claim 1 or claim 9 with a candidate agent under conditions that, in the absence of candidate agent, result in death of the cell; and

(b) subsequently measuring the ability of the candidate agent to affect the death of the cell.

22. A method for screening for a protein that regulates human mesencephalon cell death, comprising:

(a) altering the level of expression of a protein within a cell produced according to claim 1 or claim 9; and

(b) subsequently measuring the effect of the alteration on the death of the cell, and therefrom identifying a protein that regulates human mesencephalon neural precursor cell death.

23. A conditionally-immortalized human mesencephalon neural precursor cell produced according to the method of claim 1.

24. A cell according to claim 23, wherein the cell is present within a clonal cell line.

1/5

Proliferative




Fig. 1A

Differentiated




Fig. 1B

2/5




Fig. 2A



Fig. 2B

3/5




Fig. 2C




Fig. 2D

4/5



Fig. 3

5/5




Fig. 4

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/18403

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N5/10 A61K48/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PRASAD, K.N. ET AL.: "Establishment and characterization of immortalized clonal cell lines from fetal rat mesencephalic tissue" IN VITRO CELL. DEV. BIOL. ANIM., vol. 30A, no. 9, - 1994 pages 596-603, XP002122566 the whole document	6-8
Y	DI PORZIO, U. ET AL.: "Establishment and characterization of a neuronal cell line obtained by c-myc immortalization of mouse mesencephalic cells" SOCIETY FOR NEUROSCIENCE ABSTRACTS, vol. 18, no. 1-2, 1992, page P410 XP002122567 see the abstract	1-24

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 November 1999

Date of mailing of the international search report

03/12/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 851 epo nl,
Fax: (+31-70) 340-3018

Authorized officer

Alt, G

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/18403

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SAH, D.E.Y.: "Bipotent progenitor cell lines from the human CNS" NATURE BIOTECHNOLOGY, vol. 15, no. 6, 1997, pages 574-580, XP002049253 cited in the application the whole document	1-24
Y	WO 98 10058 A (SIGNAL PHARM INC) 12 March 1998 (1998-03-12) the whole document	1-24
Y	HOSHIMARU, M. ET AL.: "Differentiation of the immortalized adult neuronal progenitor cell line HC2S2 into neurons by regulatable suppression of the v-myc oncogene" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, vol. 93, no. 4, 1996, pages 1518-1523, XP002033326 cited in the application see Figure 1 the whole document	1-24
A	HARTIKKA, J. ET AL.: "Cyclic AMP, but not basic FGF, increases the in vitro survival of mesencephalic dopaminergic neurons and protects them from MPP(+)-induced degeneration" JOURNAL OF NEUROSCIENCE RESEARCH, vol. 32, 1992, pages 190-201, XP002092604 see page 195, first column, lines 7-12	11,12
A	ZHOU, J. ET AL.: "The response of human and rat fetal ventral mesencephalon in culture to the brain-derived neurotrophic factor treatment" BRAIN RESEARCH, vol. 656, no. 1, 1994, pages 147-156, XP002122571 the whole document	11,12
A	HELLER, A. ET AL.: "Glial-derived neurotrophic factor (GDNF) induced morphological differentiation of an immortalized monoclonal hybrid dopaminergic cell line of mesencephalic neuronal origin" BRAIN RESEARCH, vol. 725, no. 1, 1996, pages 132-136, XP002122572 the whole document	11,12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/18403

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9810058 A	12-03-1998	AU 4331597 A EP 0925357 A	26-03-1998 30-06-1999